

January 2016 to June 2023



Opioid and Stimulant Poisoning Hospitalizations

Surveillance of Opioid and Stimulant-Related Harms in Canada



Public Health
Agency of Canada

Agence de la santé
publique du Canada

Canada

To promote and protect the health of Canadians through leadership, partnership, innovation and action in public health.

—Public Health Agency of Canada

Prepared by the Substance-related Overdose and Mortality Surveillance Task Group (SOMS-TG) of the Federal, provincial, and territorial Special Advisory Committee (SAC) on the Epidemic of Opioid Overdoses

Également disponible en français sous le titre :
Hospitalisations pour intoxication aux opioïdes et aux stimulants
Surveillance des méfaits associés aux opioïdes et aux stimulants au Canada
Janvier 2016 à juin 2023

To obtain additional information, please contact:
Public Health Agency of Canada
Address Locator 0900C2
Ottawa, ON K1A 0K9
Tel.: 613-957-2991
Toll free: 1-866-225-0709
Fax: 613-941-5366
TTY: 1-800-465-7735
E-mail: publications-publications@hc-sc.gc.ca

© His Majesty the King in Right of Canada, as represented by the Minister of Health, 2023

Publication date: December 2023

This publication may be reproduced for personal or internal use only without permission provided the source is fully acknowledged.

Cat.: HP33-6E-PDF

ISSN: 2816-2331

Pub.: 230481



Table of Contents

Acknowledgments	4
Suggested Citation	5
Key Findings	6
Technical Notes	9
Definitions	9
How poisoning hospitalizations are counted	10
Limitations of the data on poisoning hospitalizations	11
General notes	11
Intention	11
Sex and age group	11
Fentanyl and fentanyl analogues	12
Co-occurring non-opioid poisonings	12
Co-occurring non-stimulant poisonings	12
Co-occurring non-stimulant/non-opioid poisonings	12
Data suppression	13
Table A. List of diagnosis codes included for opioid-related poisonings	14
Table B. List of diagnosis codes included for stimulant-related poisonings	14
Table C. List of diagnosis codes included for non-opioid poisonings co-occurring with opioid poisonings	15
Table D. List of diagnosis codes included for non-stimulant poisonings co-occurring with stimulant poisonings	16
Table E. List of diagnosis codes included for non-stimulant/non-opioid poisonings co-occurring with opioid or stimulant poisonings	17
Disclosure	18
Data Source	18
Appendix	19
Table 1a. Number and crude rate of total opioid-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)	20
Table 1b. Number and age-adjusted rate of total opioid-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)	23
Table 2a. Number and crude rate of accidental opioid-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)	26
Table 2b. Number and age-adjusted rate of accidental opioid-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)	29

Table 3a. Number and percentage of opioid-related poisoning hospitalizations in Canada by intention, 2016 to 2023 (January to June)	32
Table 3b. Summary of total opioid-related poisoning hospitalizations in Canada, 2016 to 2023 (January to June)	33
Table 3c. Summary of intentional opioid-related poisoning hospitalizations in Canada, 2016 to 2023 (January to June)	34
Table 4. Number of total opioid-related poisoning hospitalizations by quarter and province or territory, January 2016 to June 2023	35
Table 5. Number of accidental opioid-related poisoning hospitalizations by quarter and province or territory, January 2016 to June 2023	38
Table 6. Percentage of accidental opioid-related poisoning hospitalizations by sex and province or territory, 2016 to 2023 (January to June)	41
Table 7. Percentage of accidental opioid-related poisoning hospitalizations by age group and province or territory, 2016 to 2023 (January to June)	42
Table 8a. Number of total opioid-related poisoning hospitalizations in Canada by age group and sex, 2016 to 2023 (January to June)	50
Table 8b. Number of intentional opioid-related poisoning hospitalizations in Canada by age group and sex, 2016 to 2023 (January to June)	51
Table 8c. Number of accidental opioid-related poisoning hospitalizations by age group and sex, 2016 to 2023 (January to June)	52
Table 9. Percentage of accidental opioid-related poisoning hospitalizations involving fentanyl or fentanyl analogues by province or territory, 2016 to 2023 (January to June) .	53
Table 10. Percentage of accidental opioid-related poisoning hospitalizations by substances involved, in Canada, 2016 to 2023 (January to June)	54
Table 11a. Number and crude rate of total stimulant-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)	55
Table 11b. Number and age adjusted rate of total stimulant-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)	58
Table 12a. Number and crude rate of accidental stimulant-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)	61
Table 12b. Number and age-adjusted rate of accidental stimulant-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)	64
Table 13a. Number and percentage of stimulant-related poisoning hospitalizations by intention, 2016 to 2023 (January to June)	67
Table 13b. Summary of total stimulant-related poisoning hospitalizations in Canada, 2016 to 2023 (January to June)	68
Table 13c. Summary of intentional stimulant-related poisoning hospitalizations in Canada, 2016 to 2023 (January to June)	69
Table 14. Number of total stimulant-related poisoning hospitalizations by quarter and province or territory, January 2016 to June 2023	70
Table 15. Number of accidental stimulant-related poisoning hospitalizations by quarter and province or territory, January 2016 to June 2023	73
Table 16. Percentage of accidental stimulant-related poisoning hospitalizations by sex and province or territory, 2016 to 2023 (January to June)	76



Table 17. Percentage of accidental stimulant-related poisoning hospitalizations by age group and province or territory, 2016 to 2023 (January to June)	77
Table 18a. Number of total stimulant-related poisoning hospitalizations in Canada by sex and age group, 2016 to 2023 (January to June)	85
Table 18b. Number of intentional stimulant-related poisoning hospitalizations in Canada by age group and sex, 2016 to 2023 (January to June)	86
Table 18c. Number of accidental stimulant-related poisoning hospitalizations in Canada by age group and sex, 2016 to 2023 (January to June)	87
Table 19. Percentage of accidental stimulant-related poisoning hospitalizations by substances involved, in Canada, 2016 to 2023 (January to June).	88



Acknowledgments

We recognize that the data in this report may represent stories of pain, grief, and trauma. This report cannot adequately reflect the burden borne by Canadians. We acknowledge all those impacted by substance use, and those who work to save lives and reduce substance-related harms on individuals and communities.

This update would not be possible without the collaboration and dedication of provincial and territorial (PT) offices of Chief Coroners and Chief Medical Examiners as well as PT public health and health partners and Emergency Medical Services data providers. We would also like to acknowledge the Canadian Institute for Health Information (CIHI) for collecting and providing the data used for reporting opioid- and stimulant-related poisoning hospitalizations.



Suggested Citation

Federal, provincial, and territorial Special Advisory Committee on the Epidemic of Opioid Overdoses. Opioid- and Stimulant-related Harms in Canada. Ottawa: Public Health Agency of Canada; December 2023.
<https://health-infobase.canada.ca/substance-related-harms/opioids-stimulants/>



Key Findings

Context

Canada has seen substantially elevated numbers of opioid-related deaths and other harms since surveillance began in 2016 by the Public Health Agency of Canada. This public health crisis is having a tragic impact on people who use substances, their families, and communities across the country, and is shaped by a wide range of factors. The COVID-19 pandemic may have exacerbated the crisis, as several jurisdictions reported higher rates of fatal overdoses and other harms following the onset of the pandemic.

Presented here are the most recent available data on overdoses and deaths involving opioids and/or stimulants from January 2016 to June 2023 in Canada, where available. Recognizing that harms related to opioids, stimulants, and other substances extend beyond overdoses (poisonings) and deaths, we continue to work with federal, provincial and territorial partners to build a broad understanding of harms and substances involved to better respond to this public health crisis. For more details on additional studies and other products related to the overdose crisis, including a statement from the co-chairs of the Special Advisory Committee (SAC) on the Epidemic of Opioid Overdoses, refer to the [Data, surveillance and research page](#).

What: There was a total of 39,435 opioid-related and 16,863 stimulant-related poisoning hospitalizations from January 2016 to June 2023 in Canada (excluding Quebec)

- + A total of 3,036 opioid-related poisoning hospitalizations occurred so far in 2023 (January-June). This is an average of 17 hospitalizations per day.
- + A total of 1,107 stimulant-related poisoning hospitalizations occurred so far in 2023 (January-June). This is an average of six hospitalizations per day.
- + The median total length of hospital stay was three days for both opioid- and stimulant-related poisoning hospitalizations so far in 2023 (January-June).
- + The number of opioid-related poisoning hospitalizations in Canada reported so far in 2023 (January-June) was 11% higher compared to the same period in 2022. This is based on provisional data subject to change and should be interpreted with caution.

Data on opioid- and stimulant-related poisoning hospitalizations are not mutually exclusive. A high proportion of poisoning hospitalizations involving a stimulant also involved an opioid. Adding up those values would result in an overestimation of the burden of opioids and stimulants.



Where: A majority of accidental poisoning hospitalizations occurred in Ontario, British Columbia, and Alberta

- + So far in 2023 (January-June), most of the accidental opioid-related poisoning hospitalizations (90%) and accidental stimulant-related poisoning hospitalizations (91%) in Canada occurred in Ontario, British Columbia, and Alberta.
- + So far in 2023 (January-June), rates of accidental opioid-related hospitalizations in Canada were highest in British Columbia, Alberta, and Saskatchewan, while rates of accidental stimulant-related hospitalizations in Canada were highest in British Columbia, Saskatchewan, and Alberta.

Who: Most accidental poisoning hospitalizations were among males, with varying age distributions

- + So far in 2023 (January-June), males accounted for a higher proportion of accidental opioid-related poisoning hospitalizations (63%) and accidental stimulant-related poisoning hospitalizations (69%) than females.
- + So far in 2023 (January-June), the highest proportion of accidental opioid-related poisoning hospitalizations were among individuals aged 30 to 39 years (24%), followed by those aged 60 years or more (23%). The highest proportion of accidental stimulant-related poisoning hospitalizations were among individuals aged 30 to 39 years (28%), followed by those aged 40 to 49 years (23%).

Why: Toxicity of supply is contributing to the crisis

- + For opioid-related poisoning hospitalizations, fentanyl and its analogues were involved in 35% of accidental opioid-related poisoning hospitalizations, compared to 19% of intentional opioid-related poisoning hospitalizations so far in 2023 (January-June). The percentage of all opioid-related poisoning hospitalizations that involved fentanyl and its analogues has increased 100% since 2018 when national surveillance began, but appears to have stabilized in recent years.
- + For stimulant-related poisoning hospitalizations, fentanyl and its analogues were involved in 38% of accidental stimulant-related poisoning hospitalizations, compared to 8% of intentional stimulant-related poisoning hospitalizations so far in 2023 (January-June). The percentage of all stimulant-related poisoning hospitalizations that involved fentanyl and its analogues has increased 154% since 2018 when national surveillance began, but appears to have stabilized in recent years.



Multi-drug toxicity is a contributing factor to the crisis

- + Among accidental opioid-related poisoning hospitalizations so far in 2023 (January-June), 18% involved co-poisoning with a stimulant.
- + Among accidental stimulant-related poisoning hospitalizations so far in 2023 (January-June), 59% involved co-poisoning with an opioid.



This update is based on data submitted to or extracted by the Public Health Agency of Canada on or before November 1, 2023. Due to differences in identifying and reporting cases, comparisons over time and between provinces and territories should be interpreted with caution. Refer to the [Technical Notes](#) for more information.



Technical Notes

Definitions

Opioid-related poisoning hospitalization: Acute care hospitalizations that recorded a significant diagnosis for opioid-related poisoning. Please see [Table A](#) for a list of diagnosis codes for opioid-related poisonings.

Accidental opioid-related poisoning hospitalization: An opioid-related poisoning hospitalization that is considered to be non-intentional in nature and is defined by a diagnostic "X42" ICD -10-CA code associated with any significant opioid-related poisoning codes (T40.0-T40.4 and T40.6, see [Table A](#) for more information).

Intentional opioid-related poisoning hospitalization: An opioid-related poisoning hospitalization that occurred as a result of purposely self-inflicted harm and is defined by a diagnostic "X62" ICD-10-CA code associated with any significant opioid-related poisoning codes (T40.0-T40.4 and T40.6, see [Table A](#) for more information).

Undetermined opioid-related poisoning hospitalization: An opioid-related poisoning hospitalization that is categorized by physician documentation of undetermined/unknown intent and is defined by a diagnostic "Y12" ICD-10-CA code associated with any significant opioid-related poisoning codes (T40.0-T40.4 and T40.6, see [Table A](#) for more information).

Stimulant-related poisoning hospitalization: Acute care hospitalizations that recorded a significant diagnosis for stimulant-related poisoning. Please see [Table B](#) for a list of diagnosis codes for stimulant-related poisonings.

Accidental stimulant-related poisoning hospitalization: A stimulant-related poisoning hospitalization that is considered to be non-intentional in nature and is defined by a diagnostic "X41" or "X42" ICD -10-CA code associated with any significant stimulant-related poisoning codes (T40.5 and T43.6, see [Table B](#) for more information).

Intentional stimulant-related poisoning hospitalization: A stimulant-related poisoning hospitalization that occurred as a result of purposely self-inflicted harm and is defined by a diagnostic "X61" or "X62" ICD-10-CA code associated with any significant stimulant-related poisoning codes (T40.5 and T43.6, see [Table B](#) for more information).

Undetermined stimulant-related poisoning hospitalization: A stimulant-related poisoning hospitalization that is categorized by physician documentation of undetermined/unknown intent and is defined by a diagnostic "Y11" or "Y12" ICD-10-CA code associated with any significant stimulant-related poisoning codes (T40.5 and T43.6, see [Table B](#) for more information).

How poisoning hospitalizations are counted

Data on opioid- and stimulant-related poisoning hospitalizations are extracted from the Discharge Abstract Database (DAD), a national administrative database from the Canadian Institute for Health Information (CIHI) that compiles information on hospital discharges (including deaths, sign-outs, and transfers) from acute care institutions in all provinces and territories, except Quebec. CIHI receives the data directly from acute care facilities or from their health/regional authority or ministry/department of health. Data were accessed from the SAS grid managed by the Data Coordination and Access Program at the Public Health Agency of Canada (PHAC). Only inpatient hospitalizations from acute care facilities were included in the analyses.

Opioid and stimulant- related poisonings hospitalizations were identified as follows:

- + Poisoning diagnoses were based on the International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, Canada (ICD-10-CA), which is used to code up to 25 diagnoses per hospital record, as it is the national standard for reporting morbidity. Refer to [Table A](#) for details on the ICD-10-CA codes used to identify opioid-related poisonings (T40.0-T40.4 and T40.6) and [Table B](#) for details on the ICD-10-CA codes used to identify stimulant- related poisonings (T40.5 and T43.6). Additional ICD-10-CA codes were used to classify the hospitalizations by intent.
- + Hospitalizations were included if the opioid or stimulant-related poisoning diagnosis was considered influential to the time spent and/or treatment received while in hospital, identified by diagnosis types "M" (most responsible diagnosis (MRD)), "1" (pre-admission comorbidity), "2" (post-admission comorbidity), and "W", "X", "Y" (service transfer diagnosis).
- + Hospitalizations where the diagnosis was considered questionable, as indicated by prefix code of "Q", were excluded.

Crude and age-adjusted rates are calculated using the most current population data from Statistics Canada. Age-adjusted rates used the 2016 Canadian population as a reference and direct standardization was applied.

Crude rates: summarize the situation within a region at a certain time period and have not been adjusted for existing differences by provincial and territorial age distributions (e.g., (number of hospitalizations / population) x 100,000).

Age-adjusted rates: have been adjusted for existing differences by provincial and territorial age distributions using the 2016 Canadian population as a reference. These rates assume that all regions have the same age distributions and are useful when comparing between regions and over time.

In April 2023, PHAC began leading the data analysis and preparation of this report. Prior to this, Health Canada was conducting the analysis and preparing the reports. PHAC and Health Canada access the data through two different applications, for which the data are updated at different frequencies. Values reported in June 2023 onward may be slightly different from historically reported values.



Limitations of the data on poisoning hospitalizations

General notes

1. The unit of observation for this analysis was a discharged inpatient hospitalization visit, rather than an individual patient. The time period reported is the calendar year in which a patient was discharged from the nursing unit of the reporting facility. The province or territory reflects the location of the reporting facility where the hospitalization occurred.
2. To facilitate comparisons over time as well as across jurisdictions, age-adjusted rates per 100,000 population per year were calculated using the latest population estimates from Statistics Canada.
3. Data on hospitalizations from Quebec are not included in the Discharge Abstract Database; subsequently, Quebec is excluded from these analyses.
4. Data from Yukon, Northwest Territories and Nunavut were combined due to small numbers.
5. Data collected through DAD or the ICD-10-CA codes do not specify how the opioids were obtained (e.g., illegally or through personal prescription); the level of toxicity may differ depending on the opioid (substance(s) involved, concentration, and dosage).
6. Data after March 31, 2023 are provisional and may be less complete and/or subject to change.

Intention

7. Poisonings were further examined by the documented reason for the poisoning, based on ICD-10-CA diagnostic type "9" (external cause of injury), and classified into three categories: accidental, intentional, and undetermined/unknown.
8. Poisonings are classified as accidental unless there is clear documentation of intentional self-harm or undetermined intent.
9. When there was no documentation of intent, hospitalizations were excluded from analysis by intention, but were included in the total number of cases.

Sex and age group

10. For most provinces or territories, data on the sex of the individual was based on biological characteristics or legal documentation. Data on hospitalizations where sex was categorized as "Other" were excluded from analyses by sex, but were included in the overall analyses. Data on hospitalizations where age group was categorized as "Unknown" were excluded from analyses by age group, but were included in overall analyses. Due to rounding, percentages may not add to 100%.



Fentanyl and fentanyl analogues

11. Data on accidental opioid-related poisoning hospitalizations involving fentanyl or fentanyl analogues should be interpreted with caution as data on fentanyl-related poisoning hospitalizations became available starting April 1, 2018. As a result, fentanyl-related data presented for 2018 are limited to April to December. The ICD-10-CA classification codes to describe outcomes related to fentanyl and fentanyl analogues ("T40.40 – Poisoning by fentanyl and derivatives") were used in data collection starting 2018-19 fiscal year, as they did not exist prior to that.

Co-occurring non-opioid poisonings

12. Opioid-related poisoning hospitalizations were further examined for diagnoses of poisoning from one or more co-occurring non-opioid substance in the same hospital stay. Diagnoses for non-opioid substance-related poisoning were identified using ICD-10-CA codes and included poisonings from psychoactive substances. These codes were limited to diagnoses considered influential to the time spent/treatment received in hospital and cases where the diagnosis was considered questionable, as indicated by prefix code "Q", were excluded. Co-occurring poisonings due to non-psychoactive substances (such as non-opioid analgesics, etc.) and anti-depressants were not included. Refer to [Table C](#) for details on ICD-10-CA codes used to identify non-opioid-related poisonings.

Co-occurring non-stimulant poisonings

13. Stimulant-related poisoning hospitalizations were further examined for diagnoses of poisoning from one or more co-occurring non-stimulant substance in the same hospital stay. Diagnoses for non-stimulant substance-related poisoning were identified using ICD-10-CA codes and included poisonings from other substances. These codes were limited to diagnoses considered influential to the time spent/treatment received in hospital and cases where the diagnosis was considered questionable, as indicated by prefix code "Q", were excluded. Refer to [Table D](#) for details on ICD-10-CA codes used to identify non-stimulant-related poisonings.

Co-occurring non-stimulant/non-opioid poisonings

14. Stimulant and opioid -related poisoning hospitalizations were further examined for diagnoses of poisoning from one or more co-occurring non-stimulant/non-opioid substance in the same hospital stay. Diagnoses for non-stimulant/non-opioid substance-related poisoning were identified using ICD-10-CA codes and included poisonings from other substances. These codes were limited to diagnoses considered influential to the time spent/treatment received in hospital and cases where the diagnosis was considered questionable, as indicated by prefix code "Q", were excluded. Refer to [Table E](#) for details on ICD-10-CA codes used to identify non-stimulant/non-opioid-related poisonings.



Data suppression

Counts greater than zero and less than five have been suppressed as per CIHI privacy guidelines. Suppression was also applied in instances where all data for a province or territory fell into a single category of sex or age group. Further, in situations where a single category of a mutually exclusive variable was suppressed, an additional category was suppressed in order to address privacy concerns.



Table A. List of diagnosis codes included for opioid-related poisonings

Code	Details
T40.0	Poisoning by opium
T40.1	Poisoning by heroin
T40.2	Poisoning by other opioids
T40.20*	Poisoning by codeine and derivatives
T40.21*	Poisoning by morphine
T40.22*	Poisoning by hydromorphone
T40.23*	Poisoning by oxycodone
T40.28*	Poisoning by other opioids not elsewhere classified
T40.3	Poisoning by methadone
T40.4	Poisoning by other synthetic narcotics
T40.40*	Poisoning by fentanyl and derivatives
T40.41*	Poisoning by tramadol
T40.48*	Poisoning by other synthetic narcotics not elsewhere classified
T40.6	Poisoning by other and unspecified narcotics

*Introduced as of April 1, 2018

Table B. List of diagnosis codes included for stimulant-related poisonings

Code	Details
T40.5	Poisoning by cocaine
T43.6	Poisoning by psychostimulants with abuse potential (excl. cocaine)
T43.60**	Poisoning by methamphetamine with abuse potential
T43.68**	Poisoning by other specified psychostimulants with abuse potential
T43.69**	Poisoning by unspecified psychostimulants with abuse potential

**Introduced as of April 1, 2022



Table C. List of diagnosis codes included for non-opioid poisonings co-occurring with opioid poisonings

Code	Details
T40.5	Poisoning by cocaine
T40.7	Poisoning by cannabis (derivatives)
T40.70**	Poisoning by cannabis (derivatives), edibles and ingestibles
T40.71**	Poisoning by synthetic cannabinoids
T40.79**	Poisoning by cannabis (derivatives), unspecified
T40.8	Poisoning by lysergide (LSD)
T40.9	Poisoning by other and unspecified psychodysleptics (hallucinogens)
T42.3	Poisoning by barbiturates
T42.4	Poisoning by benzodiazepines
T42.6	Poisoning by other antiepileptic and sedative-hypnotic drugs
T42.7	Poisoning by antiepileptic and sedative-hypnotic drugs, unspecified
T43.6	Poisoning by psychostimulants with abuse potential (excl. cocaine)
T43.60**	Poisoning by methamphetamine with abuse potential
T43.68**	Poisoning by other specified psychostimulants with abuse potential
T43.69**	Poisoning by unspecified psychostimulants with abuse potential
T43.8	Poisoning by other psychotropic drugs, not elsewhere classified
T43.9	Poisoning by psychotropic drug, unspecified
T44.9	Poisoning by other and unspecified drugs primarily affecting the autonomic nervous system
T51	Toxic effect of alcohol
T51.0	Toxic effect of ethanol
T51.1	Toxic effect of methanol
T51.2	Toxic effect of 2-propanol
T51.3	Toxic effect of fusel oil
T51.8	Toxic effect of other alcohols
T51.9	Toxic effect of alcohol, unspecified

**Introduced as of April 1, 2022



Table D. List of diagnosis codes included for non-stimulant poisonings co-occurring with stimulant poisonings

Code	Details
T40.0	Poisoning by opium
T40.1	Poisoning by heroin
T40.2	Poisoning by other opioids
T40.20*	Poisoning by codeine and derivatives
T40.21*	Poisoning by morphine
T40.22*	Poisoning by hydromorphone
T40.23*	Poisoning by oxycodone
T40.28*	Poisoning by other opioids not elsewhere classified
T40.3	Poisoning by methadone
T40.4	Poisoning by other synthetic narcotics
T40.40*	Poisoning by fentanyl and derivatives
T40.41*	Poisoning by tramadol
T40.48*	Poisoning by other synthetic narcotics not elsewhere classified
T40.6	Poisoning by other and unspecified narcotics
T40.7	Poisoning by cannabis (derivatives)
T40.70**	Poisoning by cannabis (derivatives), edibles and ingestibles
T40.71**	Poisoning by synthetic cannabinoids
T40.79**	Poisoning by cannabis (derivatives), unspecified
T40.8	Poisoning by lysergide (LSD)
T40.9	Poisoning by other and unspecified psychodysleptics (hallucinogens)
T42.3	Poisoning by barbiturates
T42.4	Poisoning by benzodiazepines
T42.6	Poisoning by other antiepileptic and sedative-hypnotic drugs
T42.7	Poisoning by antiepileptic and sedative-hypnotic drugs, unspecified
T43.8	Poisoning by other psychotropic drugs, not elsewhere classified
T43.9	Poisoning by psychotropic drug, unspecified
T44.9	Poisoning by other and unspecified drugs primarily affecting the autonomic nervous system
T51	Toxic effect of alcohol
T51.0	Toxic effect of ethanol
T51.1	Toxic effect of methanol



Code	Details
T51.2	Toxic effect of 2-propanol
T51.3	Toxic effect of fusel oil
T51.8	Toxic effect of other alcohols
T51.9	Toxic effect of alcohol, unspecified

*Introduced as of April 1, 2018

**Introduced as of April 1, 2022

Table E. List of diagnosis codes included for non-stimulant/non-opioid poisonings co-occurring with opioid or stimulant poisonings

Code	Details
T40.7	Poisoning by cannabis (derivatives)
T40.70**	Poisoning by cannabis (derivatives), edibles and ingestibles
T40.71**	Poisoning by synthetic cannabinoids
T40.79**	Poisoning by cannabis (derivatives), unspecified
T40.8	Poisoning by lysergide (LSD)
T40.9	Poisoning by other and unspecified psychodysleptics (hallucinogens)
T42.3	Poisoning by barbiturates
T42.4	Poisoning by benzodiazepines
T42.6	Poisoning by other antiepileptic and sedative-hypnotic drugs
T42.7	Poisoning by antiepileptic and sedative-hypnotic drugs, unspecified
T43.8	Poisoning by other psychotropic drugs, not elsewhere classified
T43.9	Poisoning by psychotropic drug, unspecified
T44.9	Poisoning by other and unspecified drugs primarily affecting the autonomic nervous system
T51	Toxic effect of alcohol
T51.0	Toxic effect of ethanol
T51.1	Toxic effect of methanol
T51.2	Toxic effect of 2-propanol
T51.3	Toxic effect of fusel oil
T51.8	Toxic effect of other alcohols
T51.9	Toxic effect of alcohol, unspecified

**Introduced as of April 1, 2022



Disclosure

Parts of this material are based on data and information compiled and provided by CIHI. However, the analyses, conclusions, opinions and statements expressed herein are those of the authors, and not necessarily those of CIHI.

Data Source

Discharge Abstract Database (DAD), Canadian Institute for Health Information, 2016 to 2023 (January-June)

Appendix



Table 1a. Number and crude rate of total opioid-related poisoning hospitalizations by province or territory, 2016 to 2023 (January to June)

Province or territory	2016		2017		2018	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	1,156	23.8	1,407	28.5	1,418	28.3
AB	930	22.2	947	22.3	922	21.5
SK	264	23.2	244	21.2	229	19.7
MB	123	9.4	161	12.1	108	8.0
ON	1,875	13.5	2,124	15.1	2,095	14.6
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	129	16.9	101	13.2	87	11.3
NS	101	10.7	83	8.7	80	8.3
PE	22	15.0	16	10.6	20	13.0
NL	56	10.6	90	17.0	67	12.7
Territories	31	25.8	30	24.6	26	21.0
Canada	4,687	16.8	5,203	18.4	5,052	17.6

Province or territory	2019		2020		2021	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	1,196	23.5	1,586	30.8	1,954	37.6
AB	709	16.3	890	20.2	1,144	25.7
SK	264	22.5	315	26.7	321	27.2
MB	84	6.1	126	9.1	122	8.8
ON	1,981	13.6	2,044	13.9	2,446	16.5
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	104	13.4	105	13.4	98	12.4
NS	79	8.1	72	7.3	71	7.2
PE	24	15.2	17	10.5	17	10.3
NL	52	9.9	64	12.3	68	13.1
Territories	21	16.8	21	16.6	31	24.1
Canada	4,514	15.5	5,240	17.8	6,272	21.2

Province or territory	2022		2023 (Jan to Jun)	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	1,868	35.1	976	36.7
AB	893	19.7	620	27.3
SK	239	20.0	144	24.1
MB	118	8.4	62	8.8
ON	2,023	13.4	1,113	14.7
QC	n/a	n/a	n/a	n/a
NB	119	14.7	30	7.4
NS	75	7.4	40	7.8
PE	15	8.8	6	7.0
NL	57	10.8	30	11.4
Territories	24	18.5	15	23.1
Canada	5,431	18.0	3,036	20.1

Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Refer to [Technical Notes](#) for more information.

Table 1b. Number and age-adjusted rate of total opioid-related poisoning hospitalizations by province or territory, 2016 to 2023
(January to June)

Province or territory	2016		2017		2018	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	1,156	23.4	1,407	28.2	1,418	28.0
AB	930	23.3	947	23.1	922	22.1
SK	264	23.9	244	21.9	229	20.4
MB	123	9.7	161	12.4	108	8.3
ON	1,875	13.5	2,124	15.1	2,095	14.6
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	129	17.0	101	12.9	87	11.7
NS	101	10.1	83	8.5	80	8.0
PE	22	14.9	16	9.8	20	12.3
NL	56	10.3	90	16.6	67	12.8
Territories	31	29.9	30	25.6	26	21.8
Canada	4,687	16.8	5,203	18.4	5,052	17.6

Province or territory	2019		2020		2021	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	1,196	23.2	1,586	30.5	1,954	37.3
AB	709	16.8	890	20.5	1,144	25.7
SK	264	23.0	315	27.7	321	28.3
MB	84	6.3	126	9.5	122	8.9
ON	1,981	13.5	2,044	13.8	2,446	16.3
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	104	13.4	105	12.7	98	12.3
NS	79	7.8	72	6.8	71	6.7
PE	24	14.1	17	10.0	17	9.5
NL	52	9.9	64	12.3	68	13.1
Territories	21	17.7	21	17.5	31	23.8
Canada	4,514	15.5	5,240	17.8	6,272	21.1

Province or territory	2022		2023 (Jan to Jun)	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	1,868	34.9	976	36.3
AB	893	20.0	620	27.2
SK	239	20.8	144	24.9
MB	118	8.7	62	8.9
ON	2,023	13.2	1,113	14.6
QC	n/a	n/a	n/a	n/a
NB	119	14.3	30	7.3
NS	75	7.1	40	7.4
PE	15	8.7	6	6.6
NL	57	10.9	30	11.0
Territories	24	18.9	15	24.6
Canada	5,431	17.9	3,036	20.0

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.
Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Totals used to calculate age-adjusted rates do not include cases with missing age.
 Refer to [Technical Notes](#) for more information.

Table 2a. Number and crude rate of accidental opioid-related poisoning hospitalizations by province or territory, 2016 to 2023
(January to June)

Province or territory	2016		2017		2018	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	663	13.6	886	18.0	988	19.7
AB	558	13.3	624	14.7	636	14.8
SK	163	14.3	146	12.7	139	12.0
MB	67	5.1	81	6.1	61	4.5
ON	1,014	7.3	1,203	8.6	1,277	8.9
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	45	5.9	21	2.7	31	4.0
NS	53	5.6	31	3.3	39	4.1
PE	11	7.5	7	4.7	10	6.5
NL	11	2.1	19	3.6	17	3.2
Territories	11	9.2	9	7.4	15	12.1
Canada	2,596	9.3	3,027	10.7	3,213	11.2

Province or territory	2019		2020		2021	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	768	15.1	1,121	21.7	1,427	27.4
AB	490	11.2	650	14.7	923	20.8
SK	145	12.4	216	18.3	227	19.2
MB	46	3.4	66	4.8	81	5.8
ON	1,247	8.6	1,388	9.4	1,640	11.1
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	38	4.9	49	6.3	51	6.5
NS	34	3.5	39	4.0	36	3.6
PE	13	8.3	10	6.2	9	5.5
NL	19	3.6	22	4.2	20	3.8
Territories	5	4.0	11	8.7	18	14.0
Canada	2,805	9.6	3,572	12.1	4,432	15.0

Province or territory	2022		2023 (Jan to Jun)	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	1,401	26.3	738	27.7
AB	679	14.9	510	22.5
SK	179	15.0	110	18.4
MB	77	5.5	41	5.8
ON	1,335	8.8	757	10.0
QC	n/a	n/a	n/a	n/a
NB	60	7.4	18	4.4
NS	30	2.9	23	4.5
PE	9	5.3	Suppr.	Suppr.
NL	18	3.4	9	3.4
Territories	13	10.0	10	15.4
Canada	3,801	12.6	2,216	14.7

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.
Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Refer to [Technical Notes](#) for more information.

Table 2b. Number and age-adjusted rate of accidental opioid-related poisoning hospitalizations by province or territory, 2016 to 2023
(January to June)

Province or territory	2016		2017		2018	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	663	13.4	886	17.7	988	19.5
AB	558	14.4	624	15.4	636	15.3
SK	163	14.7	146	12.9	139	12.3
MB	67	5.3	81	6.2	61	4.7
ON	1,014	7.3	1,203	8.5	1,277	8.9
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	45	5.6	21	2.4	31	3.8
NS	53	5.1	31	3.0	39	3.7
PE	11	7.3	7	4.6	10	5.6
NL	11	1.9	19	3.4	17	3.0
Territories	11	10.9	9	7.4	15	13.1
Canada	2,596	9.3	3,027	10.7	3,213	11.2

Province or territory	2019		2020		2021	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	768	14.9	1,121	21.5	1,427	27.3
AB	490	11.7	650	15.0	923	20.8
SK	145	12.6	216	19.1	227	20.1
MB	46	3.4	66	5.1	81	5.9
ON	1,247	8.5	1,388	9.4	1,640	10.9
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	38	4.7	49	5.6	51	6.3
NS	34	3.2	39	3.4	36	3.3
PE	13	7.0	10	5.3	9	4.7
NL	19	3.4	22	4.0	20	3.6
Territories	5	5.5	11	10.1	18	14.0
Canada	2,805	9.6	3,572	12.1	4,432	14.9

Province or territory	2022		2023 (Jan to Jun)	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	1,401	26.2	738	27.5
AB	679	15.2	510	22.3
SK	179	15.6	110	19.0
MB	77	5.7	41	5.9
ON	1,335	8.7	757	9.9
QC	n/a	n/a	n/a	n/a
NB	60	6.8	18	4.0
NS	30	2.9	23	3.9
PE	9	5.0	Suppr.	Suppr.
NL	18	3.1	9	3.1
Territories	13	10.3	10	17.8
Canada	3,801	12.5	2,216	14.6

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals used to calculate age-adjusted rates do not include cases with missing age.

Refer to [Technical Notes](#) for more information.

Table 3a. Number and percentage of opioid-related poisoning hospitalizations in Canada by intention, 2016 to 2023 (January to June)

Intention	2016			2017			2018			2019			2020			2021			2022			2023 (Jan to Jun)		
	Number	%																						
Total	4,687	100%	5,203	100%	5,052	100%	4,514	100%	5,240	100%	6,272	100%	5,431	100%	5,431	100%	3,036	100%	3,036	100%	3,036	100%		
Accidental	2,596	55%	3,027	58%	3,213	64%	2,805	62%	3,572	68%	4,432	71%	3,801	70%	2,216	73%								
Intentional	1,370	29%	1,387	27%	1,247	25%	1,170	26%	1,067	20%	1,159	18%	1,089	20%	521	17%								
Unknown	600	13%	651	13%	561	11%	530	12%	596	11%	679	11%	543	10%	296	10%								

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

The total number includes opioid-related poisoning hospitalizations for which the intent was not recorded.

Intention categories may sum to greater than the total number due to multiple intentions being recorded on the same record.

Refer to [Technical Notes](#) for more information.

Table 3b. Summary of total opioid-related poisoning hospitalizations in Canada, 2016 to 2023 (January to June)

		2016	2017	2018	2019	2020	2021	2022	2023 (Jan to Jun)
Total	Number	4,687	5,203	5,052	4,514	5,240	6,272	5,431	3,036
	Crude rate	16.8	18.4	17.6	15.5	17.8	21.2	18.0	20.1
	Age adjusted rate	16.8	18.4	17.6	15.5	17.8	21.1	17.9	20.0
Percent by sex	Male	50%	55%	55%	54%	60%	62%	60%	61%
	Female	50%	45%	45%	46%	40%	38%	40%	39%
	0 to 19	6%	6%	7%	6%	5%	5%	5%	5%
	20 to 29	17%	18%	18%	17%	17%	17%	15%	15%
	30 to 39	16%	19%	19%	20%	21%	23%	21%	23%
	40 to 49	15%	14%	16%	14%	16%	18%	17%	18%
	50 to 59	20%	19%	18%	18%	18%	16%	17%	15%
60 or more		27%	25%	23%	25%	22%	22%	24%	23%
	Fentanyl or fentanyl analogues	n/a	n/a	16%	21%	28%	32%	31%	32%
	Co-occurring non-opioid poisonings	36%	36%	35%	35%	33%	28%	29%	28%
	Co-occurring stimulant poisonings	16%	18%	20%	19%	20%	17%	17%	18%
Percent involving categories not mutually exclusive	Co-occurring non-stimulant/non-opioid poisonings	28%	25%	23%	22%	20%	17%	19%	18%
	Median total length of stay (in days)	3	3	3	3	3	3	3	3
	Not available (n/a)								

Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded. Totals used to calculate age-adjusted rates do not include cases with missing age.

The ICD-10-CA code for poisoning by fentanyl and derivatives was introduced April 2018. Fentanyl-specific data were not available prior to this date.

Data on opioid-related and stimulant-related poisonings hospitalizations are not mutually exclusive. A proportion of poisoning hospitalizations involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

Refer to the [Technical Notes](#) for more information.

Table 3c. Summary of intentional opioid-related poisoning hospitalizations in Canada, 2016 to 2023 (January to June)

		2016	2017	2018	2019	2020	2021	2022	2023 (Jan to Jun)
Total	Number	1,370	1,387	1,247	1,170	1,067	1,159	1,089	521
Crude rate	4.9	4.9	4.4	4.0	3.6	3.9	3.6	3.6	3.4
Age adjusted rate	4.9	4.9	4.4	4.0	3.6	3.9	3.6	3.6	3.4
Percent by sex	Male	43%	46%	41%	41%	45%	46%	45%	48%
	Female	57%	54%	59%	59%	55%	54%	55%	52%
Percent by age group	0 to 19	9%	10%	12%	11%	9%	10%	9%	11%
	20 to 29	18%	19%	18%	18%	20%	17%	18%	17%
	30 to 39	17%	19%	17%	18%	19%	19%	16%	18%
	40 to 49	18%	16%	17%	17%	16%	16%	16%	18%
	50 to 59	22%	20%	18%	18%	16%	15%	16%	14%
	60 or more	16%	16%	18%	18%	19%	22%	25%	23%
	Fentanyl or fentanyl analogues	n/a	n/a	7%	11%	14%	18%	19%	19%
Percent involving (categories not mutually exclusive)	Co-occurring non-opioid poisonings	51%	49%	48%	50%	48%	43%	44%	44%
	Co-occurring stimulant poisonings	16%	17%	18%	20%	20%	16%	16%	17%
	Co-occurring non-stimulant/non-opioid poisonings	44%	41%	39%	39%	36%	34%	37%	34%

Not available (n/a)

Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals used to calculate age-adjusted rates do not include cases with missing age.

The ICD-10-CA code for poisoning by fentanyl and derivatives was introduced April 2018. Fentanyl-specific data were not available prior to this date.

Data on opioid-related and stimulant-related poisoning hospitalizations are not mutually exclusive. A proportion of poisoning hospitalizations involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.
Refer to the [Technical Notes](#) for more information.

Table 4. Number of total opioid-related poisoning hospitalizations by quarter and province or territory, January 2016 to June 2023

Province or territory	2016				2017				2018			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	304	269	278	305	338	375	358	336	293	382	380	363
AB	229	239	239	223	235	235	256	221	202	229	271	220
SK	75	59	72	58	55	60	64	65	52	65	61	51
MB	25	35	26	37	40	49	32	40	30	27	23	28
ON	402	500	458	515	493	537	578	516	481	566	519	529
QC	n/a											
NB	24	26	46	33	29	28	19	25	24	22	22	19
NS	18	23	32	28	28	17	18	20	13	33	16	18
PE	Suppr.	Suppr.	7	6	7	Suppr.	Suppr.	6	7	Suppr.	7	Suppr.
NL	15	16	14	11	21	24	28	17	19	20	19	9
Territories	Suppr.	13	Suppr.	8	10	8	6	6	11	Suppr.	8	Suppr.
Canada	1,092	1,180	1,172	1,224	1,256	1,333	1,359	1,252	1,132	1,344	1,326	1,237

Province or territory	2019				2020				2021			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	315	308	314	259	281	379	452	474	461	474	497	522
AB	171	174	214	150	188	226	235	241	210	272	355	307
SK	56	73	67	68	50	83	87	95	72	79	77	93
MB	18	17	21	28	26	30	41	29	31	26	33	32
ON	515	573	454	439	455	519	543	527	595	618	619	614
QC	n/a											
NB	30	20	27	27	23	28	29	25	27	26	25	20
NS	23	19	18	19	15	22	13	22	18	20	14	19
PE	Suppr.	8	6	Suppr.	6	Suppr.	Suppr.	7	5	Suppr.	5	Suppr.
NL	9	16	19	8	16	22	12	14	11	17	24	16
Territories	Suppr.	8	Suppr.	5	7	6	Suppr.	Suppr.	Suppr.	Suppr.	8	12
Canada	1,137	1,216	1,140	1,003	1,067	1,315	1,412	1,434	1,430	1,532	1,657	1,635

Province or territory	2022				2023	
	Q1	Q2	Q3	Q4	Q1	Q2
BC	446	491	468	463	529	447
AB	267	224	203	199	287	333
SK	65	55	64	55	64	80
MB	31	33	25	29	34	28
ON	485	496	528	514	541	572
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	23	35	35	26	20	10
NS	18	20	13	24	21	19
PE	Suppr.	6	Suppr.	Suppr.	Suppr.	Suppr.
NL	15	9	19	14	21	9
Territories	Suppr.	9	8	Suppr.	10	5
Canada	1,350	1,378	1,363	1,324	1,527	1,503

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Q1 = January to March; Q2 = April to June; Q3 = July to September; Q4 = October to December

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Totals for Canada by year may not match those from other tables, as they do not include the suppressed numbers in provinces/territories, in order to comply with privacy and confidentiality rules.
 Refer to [Technical Notes](#) for more information.

Table 5. Number of accidental opioid-related poisoning hospitalizations by quarter and province or territory, January 2016 to June 2023

Province or territory	2016				2017				2018			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	164	147	155	197	204	236	221	225	190	280	266	252
AB	135	140	156	127	151	146	169	158	138	165	168	165
SK	42	37	51	33	28	36	42	40	31	36	37	35
MB	12	17	15	23	16	27	18	20	16	15	12	18
ON	220	267	243	284	256	307	356	284	269	370	327	311
QC	n/a											
NB	7	13	15	10	Suppr.	Suppr.	6	6	8	9	8	6
NS	7	11	22	13	10	5	8	8	Suppr.	20	Suppr.	8
PE	Suppr.	Suppr.	Suppr.	0	0	Suppr.						
NL	Suppr.	5	Suppr.	Suppr.	5	6	Suppr.	Suppr.	Suppr.	7	Suppr.	Suppr.
Territories	Suppr.	7	Suppr.	5	Suppr.							
Canada	587	637	657	687	670	763	820	741	659	902	823	795

Province or territory	2019				2020				2021			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	212	204	190	162	177	258	327	359	317	355	375	380
AB	114	129	149	98	124	164	168	194	160	225	282	256
SK	22	42	39	42	30	58	58	70	52	46	55	74
MB	9	12	10	15	11	13	25	17	19	16	24	22
ON	336	363	269	279	310	364	379	335	386	423	411	420
QC	n/a	n/a	n/a									
NB	8	6	11	13	11	14	13	11	12	15	15	9
NS	8	5	9	12	8	12	7	12	7	14	6	9
PE	Suppr.	6	Suppr.	Suppr.	0							
NL	6	Suppr.	6	Suppr.	6	13	Suppr.	Suppr.	Suppr.	6	5	Suppr.
Territories	Suppr.	Suppr.	0	Suppr.	5	Suppr.	Suppr.	Suppr.	5	0	7	6
Canada	715	767	683	621	682	896	977	998	958	1,100	1,180	1,176

Province or territory	2022				2023	
	Q1	Q2	Q3	Q4	Q1	Q2
BC	347	370	353	331	398	340
AB	209	169	148	153	228	282
SK	50	41	47	41	47	63
MB	18	24	16	19	23	18
ON	323	327	351	334	361	396
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	12	22	14	12	12	6
NS	10	6	5	9	11	12
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	5	7	Suppr.	Suppr.
Territories	Suppr.	6	5	Suppr.	Suppr.	Suppr.
Canada	969	965	944	906	1,080	1,117

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Q1 = January to March; Q2 = April to June; Q3 = July to September; Q4 = October to December

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Totals for Canada by year may not match those from other tables, as they do not include the suppressed numbers in provinces/territories, in order to comply with privacy and confidentiality rules.
 Refer to [Technical Notes](#) for more information.

Table 6. Percentage of accidental opioid-related poisoning hospitalizations by sex and province or territory, 2016 to 2023
(January to June)

Province or territory	2016		2017		2018		2019		2020		2021		2022		2023 (Jan to Jun)	
	Female	Male	Female	Male												
BC	37%	63%	31%	69%	32%	68%	34%	66%	30%	70%	30%	70%	29%	71%	31%	69%
AB	50%	50%	43%	57%	42%	58%	46%	54%	38%	62%	37%	63%	38%	62%	38%	62%
SK	50%	50%	49%	51%	44%	56%	44%	52%	48%	47%	53%	40%	60%	47%	53%	45%
MB	52%	48%	43%	57%	59%	41%	54%	46%	39%	61%	36%	64%	44%	56%	56%	44%
ON	51%	49%	46%	54%	44%	56%	43%	57%	38%	62%	36%	64%	37%	63%	38%	62%
QC	n/a	n/a														
NB	36%	64%	71%	29%	55%	45%	42%	58%	49%	51%	37%	63%	60%	40%	56%	44%
NS	58%	42%	45%	55%	41%	59%	53%	47%	62%	38%	47%	53%	60%	40%	48%	52%
PE	45%	55%	Suppr.	Suppr.												
NL	Suppr.	Suppr.	32%	68%	59%	41%	47%	53%	50%	50%	45%	55%	Suppr.	Suppr.	Suppr.	Suppr.
Territories	Suppr.	38%	62%	50%	50%											
Canada	47%	53%	41%	59%	41%	59%	42%	58%	37%	63%	35%	65%	35%	65%	37%	63%

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.
Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
The data do not include cases with missing sex, or other sex.

Totals used to calculate Canada proportions do not include the suppressed numbers, in order to comply with privacy and confidentiality rules.
Refer to [Technical Notes](#) for more information.

Table 7. Percentage of accidental opioid-related poisoning hospitalizations by age group and province or territory, 2016 to 2023 (January to June)

2016

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	3%	20%	21%	16%	15%	24%
AB	4%	16%	12%	16%	18%	33%
SK	9%	14%	19%	14%	20%	24%
MB	Suppr.	12%	Suppr.	19%	19%	36%
ON	4%	13%	14%	11%	20%	39%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	13%	Suppr.	22%	22%	36%
NS	Suppr.	11%	Suppr.	Suppr.	15%	60%
PE	0%	Suppr.	0%	0%	Suppr.	55%
NL	Suppr.	0%	Suppr.	Suppr.	Suppr.	45%
Territories	Suppr.	Suppr.	Suppr.	0%	Suppr.	Suppr.
Canada	4%	16%	15%	14%	19%	34%

2017

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	5%	22%	23%	17%	15%	18%
AB	3%	19%	21%	13%	18%	27%
SK	12%	17%	17%	15%	17%	22%
MB	9%	25%	21%	7%	12%	26%
ON	4%	14%	16%	13%	20%	34%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	0%	Suppr.	0%	Suppr.	24%	57%
NS	Suppr.	Suppr.	Suppr.	Suppr.	19%	55%
PE	0%	Suppr.	Suppr.	0%	0%	71%
NL	0%	Suppr.	Suppr.	Suppr.	37%	32%
Territories	0%	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	5%	17%	19%	14%	18%	28%

2018

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	5%	22%	22%	17%	17%	17%
AB	4%	21%	18%	16%	17%	25%
SK	11%	18%	19%	14%	16%	22%
MB	8%	23%	11%	10%	25%	23%
ON	3%	13%	20%	14%	18%	32%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	Suppr.	Suppr.	Suppr.	19%	45%
NS	Suppr.	Suppr.	Suppr.	Suppr.	18%	51%
PE	0%	0%	0%	Suppr.	0%	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	47%
Territories	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	33%
Canada	4%	18%	19%	15%	17%	26%

2019

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	7%	18%	24%	14%	17%	20%
AB	4%	14%	20%	12%	16%	35%
SK	15%	22%	15%	15%	13%	19%
MB	Suppr.	22%	20%	Suppr.	13%	28%
ON	3%	16%	20%	13%	18%	31%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	Suppr.	Suppr.	13%	21%	45%
NS	Suppr.	Suppr.	Suppr.	21%	Suppr.	53%
PE	0%	0%	Suppr.	Suppr.	Suppr.	77%
NL	0%	Suppr.	Suppr.	Suppr.	42%	32%
Territories	0%	Suppr.	0%	Suppr.	0%	Suppr.
Canada	5%	16%	20%	13%	17%	29%

2020

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	4%	18%	20%	19%	18%	20%
AB	4%	16%	22%	16%	18%	24%
SK	4%	14%	26%	19%	18%	19%
MB	Suppr.	17%	Suppr.	14%	26%	29%
ON	5%	15%	21%	14%	18%	26%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	Suppr.	18%	Suppr.	Suppr.	61%
NS	Suppr.	Suppr.	Suppr.	Suppr.	18%	64%
PE	0%	Suppr.	Suppr.	0%	Suppr.	60%
NL	Suppr.	Suppr.	Suppr.	Suppr.	41%	32%
Territories	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	4%	16%	21%	16%	18%	25%

2021

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	4%	15%	22%	21%	20%	18%
AB	3%	19%	26%	18%	13%	20%
SK	6%	17%	28%	17%	19%	13%
MB	Suppr.	38%	26%	Suppr.	11%	12%
ON	3%	15%	23%	17%	15%	27%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	10%	24%	18%	Suppr.	39%
NS	Suppr.	Suppr.	19%	Suppr.	14%	47%
PE	0%	Suppr.	Suppr.	0%	Suppr.	67%
NL	Suppr.	Suppr.	Suppr.	Suppr.	35%	35%
Territories	Suppr.	Suppr.	28%	Suppr.	Suppr.	Suppr.
Canada	4%	16%	24%	18%	16%	22%

2022

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	4%	14%	25%	19%	19%	19%
AB	5%	18%	23%	16%	14%	24%
SK	7%	15%	22%	18%	17%	21%
MB	Suppr.	Suppr.	23%	17%	16%	23%
ON	4%	12%	20%	16%	17%	31%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	18%	Suppr.	Suppr.	17%	55%
NS	Suppr.	Suppr.	Suppr.	20%	17%	33%
PE	0%	0%	Suppr.	Suppr.	Suppr.	56%
NL	0%	Suppr.	Suppr.	Suppr.	39%	33%
Territories	0%	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	4%	14%	22%	17%	17%	25%

2023 (Jan to Jun)

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	4%	17%	26%	18%	18%	18%
AB	6%	16%	27%	20%	12%	17%
SK	5%	19%	21%	18%	9%	27%
MB	Suppr.	22%	41%	12%	12%	Suppr.
ON	3%	12%	22%	19%	15%	29%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	0%	Suppr.	Suppr.	Suppr.	Suppr.	56%
NS	Suppr.	0%	Suppr.	0%	Suppr.	74%
PE	Suppr.	0%	0%	0%	0%	Suppr.
NL	Suppr.	0%	0%	Suppr.	Suppr.	Suppr.
Territories	0%	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	4%	15%	24%	18%	15%	23%

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

The data do not include cases with missing age.

Totals used to calculate Canada proportions do not include the suppressed numbers, in order to comply with privacy and confidentiality rules.

Refer to [Technical Notes](#) for more information.

Table 8a. Number of total opioid-related poisoning hospitalizations in Canada by age group and sex, 2016 to 2023 (January to June)

Age group by sex	2016		2017		2018		2019		2020		2021		2022		2023 (Jan to Jun)	
	Female	Male	Female	Male												
0 to 19	151	111	186	117	207	123	149	128	157	127	178	138	164	119	85	80
20 to 29	289	490	318	620	335	571	289	476	305	565	343	712	287	539	188	279
30 to 39	295	450	325	641	309	641	346	552	363	744	505	922	381	773	243	452
40 to 49	371	341	316	437	345	457	278	370	320	541	386	738	306	613	197	352
50 to 59	467	466	494	471	445	441	404	401	361	581	352	640	357	568	155	297
60 or more	754	499	678	598	644	533	614	507	595	578	627	720	658	660	306	389

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals for sex by age group may not match those from other tables as they do not include cases with missing sex, other sex or missing age.

Refer to [Technical Notes](#) for more information.

Table 8b. Number of intentional opioid-related poisoning hospitalizations in Canada by age group and sex, 2016 to 2023 (January to June)

Age group by sex	2016		2017		2018		2019		2020		2021		2022		2023 (Jan to Jun)	
	Female	Male	Female	Male												
0 to 19	93	37	107	24	113	41	91	33	78	22	92	28	78	23	41	14
20 to 29	124	121	127	136	128	96	122	93	111	103	89	107	89	103	46	43
30 to 39	124	115	117	142	102	110	123	86	107	100	108	116	87	90	54	38
40 to 49	154	88	113	114	121	93	100	100	89	82	113	73	91	81	39	53
50 to 59	164	136	172	112	145	75	132	84	92	84	86	83	100	78	37	34
60 or more	128	85	114	108	121	102	119	87	111	87	130	130	150	118	56	66

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals for sex by age group may not match those from other tables as they do not include cases with missing sex, other sex or missing age.
Refer to [Technical Notes](#) for more information.

Table 8c. Number of accidental opioid-related poisoning hospitalizations by age group and sex, 2016 to 2023 (January to June)

Age group by sex	2016		2017		2018		2019		2020		2021		2022		2023 (Jan to Jun)	
	Female	Male	Female	Male												
0 to 19	49	62	64	76	77	69	54	85	68	95	77	94	75	88	37	62
20 to 29	120	282	154	378	169	398	135	321	166	395	214	516	164	373	131	201
30 to 39	141	247	156	410	173	454	181	383	206	536	345	693	255	594	169	366
40 to 49	157	198	160	260	184	299	137	234	193	384	238	571	182	465	140	262
50 to 59	236	240	252	282	255	299	230	240	239	413	228	487	220	437	103	230
60 or more	517	345	460	374	460	375	436	432	369	443	444	519	461	484	227	280

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals for sex by age group may not match those from other tables as they do not include cases with missing sex, other sex or missing age.

Refer to [Technical Notes](#) for more information.

Table 9. Percentage of accidental opioid-related poisoning hospitalizations involving fentanyl or fentanyl analogues by province or territory, 2016 to 2023 (January to June)

Province or territory	2016	2017	2018 (Apr to Dec)	2019	2020	2021	2022	2023 (Jan to Jun)
BC	n/a	n/a	33%	40%	42%	41%	43%	39%
AB	n/a	n/a	16%	20%	19%	22%	25%	27%
SK	n/a	n/a	7%	6%	25%	30%	31%	40%
MB	n/a	n/a	11%	13%	14%	30%	17%	20%
ON	n/a	n/a	13%	23%	33%	39%	32%	37%
QC	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
NB	n/a	n/a	0%	Suppr.	Suppr.	Suppr.	17%	Suppr.
NS	n/a	n/a	0%	0%	0%	Suppr.	0%	0%
PE	n/a	n/a	Suppr.	0%	0%	0%	0%	0%
NL	n/a	n/a	Suppr.	0%	0%	Suppr.	0%	0%
Territories	n/a	n/a	40%	Suppr.	45%	Suppr.	54%	70%
Canada	n/a	n/a	19%	25%	32%	35%	34%	35%

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

The ICD-10-CA code for poisoning by fentanyl and derivatives was introduced April 2018. Fentanyl-specific data were not available prior to this date.

Refer to [Technical Notes](#) for more information.

Table 10. Percentage of accidental opioid-related poisoning hospitalizations by substances involved, in Canada, 2016 to 2023
(January to June)

	2016	2017	2018	2019	2020	2021	2022	2023 (Jan to Jun)
Total number	2,596	3,027	3,213	2,805	3,572	4,432	3,801	2,216
Percent involving co-occurring non-opioid poisonings	29%	30%	30%	28%	29%	24%	25%	25%
Percent involving co-occurring stimulant poisonings	16%	18%	20%	18%	20%	17%	17%	18%
Percent involving co-occurring non-stimulant/non-opioid poisonings	20%	19%	18%	17%	16%	13%	15%	15%

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Data on opioid-related and stimulant-related poisoning hospitalizations are not mutually exclusive. A proportion of poisoning hospitalizations involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.
Refer to [Technical Notes](#) for more information.

Table 11a. Number and crude rate of total stimulant-related poisoning hospitalizations by province or territory, 2016 to 2023
(January to June)

Province or territory	2016		2017		2018	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	480	9.9	583	11.8	548	10.9
AB	370	8.8	401	9.5	399	9.3
SK	120	10.6	108	9.4	126	10.8
MB	66	5.0	82	6.1	64	4.7
ON	722	5.2	916	6.5	1,107	7.7
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	51	6.7	54	7.0	53	6.9
NS	32	3.4	37	3.9	27	2.8
PE	5	3.4	9	6.0	6	3.9
NL	28	5.3	33	6.2	31	5.9
Territories	11	9.2	12	9.8	12	9.7
Canada	1,885	6.8	2,235	7.9	2,373	8.3

Province or territory	2019		2020		2021	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	524	10.3	606	11.8	663	12.7
AB	321	7.4	425	9.6	402	9.0
SK	159	13.6	143	12.1	162	13.7
MB	58	4.2	56	4.1	54	3.9
ON	975	6.7	1,111	7.5	1,002	6.8
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	56	7.2	59	7.5	60	7.6
NS	32	3.3	46	4.7	34	3.4
PE	6	3.8	7	4.3	Suppr.	Suppr.
NL	36	6.9	43	8.2	32	6.1
Territories	12	9.6	20	15.8	13	10.1
Canada	2,179	7.5	2,516	8.5	2,422	8.2

Province or territory	2022		2023 (Jan to Jun)	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	673	12.7	305	11.5
AB	306	6.7	168	7.4
SK	116	9.7	75	12.6
MB	54	3.8	22	3.1
ON	844	5.6	482	6.4
QC	n/a	n/a	n/a	n/a
NB	61	7.5	19	4.7
NS	36	3.5	15	2.9
PE	Suppr.	Suppr.	Suppr.	Suppr.
NL	37	7.0	8	3.0
Territories	19	14.6	13	20.0
Canada	2,146	7.1	1,107	7.3

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Refer to [Technical Notes](#) for more information.

Table 11b. Number and age adjusted rate of total stimulant-related poisoning hospitalizations by province or territory, 2016 to 2018
(January to June)

Province or territory	2016		2017		2018	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	480	9.9	583	11.9	548	11.0
AB	370	8.5	401	9.1	399	9.0
SK	120	10.5	108	9.4	126	11.0
MB	66	5.1	82	6.2	64	4.7
ON	722	5.2	916	6.5	1,107	7.8
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	51	7.2	54	7.9	53	7.7
NS	32	3.7	37	4.2	27	3.0
PE	5	3.8	9	6.6	6	4.3
NL	28	6.0	33	7.0	31	6.9
Territories	11	9.5	12	8.9	12	8.9
Canada	1,885	6.8	2,235	7.9	2,373	8.3

Province or territory	2019		2020		2021	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	524	10.4	606	11.8	663	13.0
AB	321	7.2	425	9.4	402	8.9
SK	159	13.5	143	12.3	162	13.9
MB	58	4.2	56	4.1	54	3.8
ON	975	6.7	1,111	7.6	1,002	6.8
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	56	8.0	59	8.3	60	8.3
NS	32	3.5	46	5.0	34	3.8
PE	6	4.1	7	4.4	Suppr.	Suppr.
NL	36	7.8	43	9.3	32	7.0
Territories	12	8.7	20	14.8	13	8.9
Canada	2,179	7.5	2,516	8.6	2,422	8.3

Province or territory	Number	2022		2023 (Jan to Jun)	
		Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number
BC	673	12.8	305	11.7	
AB	306	6.6	168	7.3	
SK	116	9.9	75	12.7	
MB	54	4.0	22	3.2	
ON	844	5.6	482	6.5	
QC	n/a	n/a	n/a	n/a	
NB	61	8.3	19	5.2	
NS	36	3.7	15	3.1	
PE	Suppr.	Suppr.	Suppr.	Suppr.	
NL	37	8.2	8	3.2	
Territories	19	13.6	13	19.3	
Canada	2,146	7.2	1,107	7.4	

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.
Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Totals used to calculate age-adjusted rates do not include cases with missing age.
 Refer to [Technical Notes](#) for more information.

Table 12a. Number and crude rate of accidental stimulant-related poisoning hospitalizations by province or territory, 2016 to 2023
(January to June)

Province or territory	2016		2017		2018	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	295	6.1	342	6.9	378	7.5
AB	191	4.6	228	5.4	223	5.2
SK	60	5.3	58	5.0	69	5.9
MB	33	2.5	46	3.4	27	2.0
ON	333	2.4	457	3.2	583	4.1
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	11	1.4	5	0.7	5	0.6
NS	9	1.0	9	0.9	7	0.7
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	6	1.1
Territories	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	932	3.4	1,145	4.1	1,298	4.6

Province or territory	2019		2020		2021	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	288	5.7	375	7.3	425	8.2
AB	188	4.3	261	5.9	246	5.5
SK	63	5.4	72	6.1	78	6.6
MB	28	2.0	25	1.8	32	2.3
ON	550	3.8	670	4.5	576	3.9
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	14	1.8	27	3.4	20	2.5
NS	10	1.0	14	1.4	14	1.4
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	6	1.1	5	1.0	5	1.0
Territories	Suppr.	Suppr.	Suppr.	Suppr.	6	4.7
Canada	1,147	4.0	1,449	4.9	1,402	4.7

Province or territory	2022		2023 (Jan to Jun)	
	Number	Crude rate per 100,000 population	Number	Crude rate per 100,000 population
BC	440	8.3	199	7.5
AB	169	3.7	101	4.4
SK	56	4.7	38	6.4
MB	24	1.7	9	1.3
ON	486	3.2	298	3.9
QC	n/a	n/a	n/a	n/a
NB	21	2.6	7	1.7
NS	10	1.0	5	1.0
PE	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.
Territories	10	77	Suppr.	Suppr.
Canada	1,216	4.0	657	4.4

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.
Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Refer to [Technical Notes](#) for more information.

Table 12b. Number and age-adjusted rate of accidental stimulant-related poisoning hospitalizations by province or territory, 2016 to 2023
(January to June)

Province or territory	2016		2017		2018	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	295	6.1	342	7.0	378	7.6
AB	191	4.4	228	5.2	223	5.0
SK	60	5.2	58	5.0	69	6.1
MB	33	2.5	46	3.5	27	2.0
ON	333	2.4	457	3.3	583	4.1
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	11	1.5	5	0.8	5	0.7
NS	9	1.0	9	1.0	7	0.8
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	Suppr.	Suppr.	Suppr.	Suppr.	6	1.3
Territories	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Canada	932	3.4	1,145	4.1	1,298	4.6

Province or territory	2019		2020		2021	
	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population
BC	288	5.7	375	7.3	425	8.2
AB	188	4.2	261	5.8	246	5.4
SK	63	5.4	72	6.3	78	6.7
MB	28	2.0	25	1.9	32	2.4
ON	550	3.8	670	4.6	576	3.9
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	14	2.0	27	3.7	20	2.7
NS	10	1.1	14	1.5	14	1.5
PE	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	6	1.4	5	1.2	5	1.1
Territories	Suppr.	Suppr.	Suppr.	Suppr.	6	4.0
Canada	1,147	4.0	1,449	5.0	1,402	4.8

Province or territory	Number	2022		2023 (Jan to Jun)	
		Age-adjusted rate per 100,000 population	Number	Age-adjusted rate per 100,000 population	Number
BC	440	8.3	199	7.5	
AB	169	3.6	101	4.4	
SK	56	4.9	38	6.5	
MB	24	1.8	9	1.3	
ON	486	3.3	298	4.0	
QC	n/a	n/a	n/a	n/a	
NB	21	2.9	7	1.9	
NS	10	1.0	5	1.0	
PE	Suppr.	Suppr.	Suppr.	Suppr.	
NL	Suppr.	Suppr.	Suppr.	Suppr.	
Territories	10	7.3	Suppr.	Suppr.	
Canada	1,216	4.1	657	4.5	

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals used to calculate age-adjusted rates do not include cases with missing age.

Refer to [Technical Notes](#) for more information.

Table 13a. Number and percentage of stimulant-related poisoning hospitalizations by intention, 2016 to 2023 (January to June)

Intention	2016			2017			2018			2019			2020			2021			2022			2023 (Jan to Jun)		
	Number	%	Number	%																				
Total	1,885	100%	2,235	100%	2,373	100%	2,179	100%	2,516	100%	2,422	100%	2,146	100%	2,107	100%	2,146	100%	2,107	100%	2,146	100%	2,107	100%
Accidental	932	49%	1,145	51%	1,298	55%	1,147	53%	1,449	58%	1,402	58%	1,216	57%	657	59%	657	57%	657	57%	657	57%	657	59%
Intentional	649	34%	699	31%	696	29%	703	32%	729	29%	698	29%	657	31%	329	30%	329	31%	329	31%	329	30%	329	30%
Unknown	288	15%	382	17%	376	16%	329	15%	327	13%	334	14%	281	13%	122	11%	122	11%	122	11%	122	11%	122	11%

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

The total number includes stimulant-related poisoning hospitalizations for which the intent was not recorded.

Intention categories may sum to greater than the total number due to multiple intentions recorded on the same record.

Refer to [Technical Notes](#) for more information.

Table 13b. Summary of total stimulant-related poisoning hospitalizations in Canada, 2016 to 2023 (January to June)

		2016	2017	2018	2019	2020	2021	2022	2023 (Jan to Jun)
Total	Number	1,885	2,235	2,373	2,179	2,516	2,422	2,146	1,107
	Crude rate	6.8	7.9	8.3	7.5	8.5	8.2	7.1	7.3
	Age adjusted rate	6.8	7.9	8.3	7.5	8.6	8.3	7.2	7.4
Percent by sex	Male	60%	62%	61%	60%	61%	60%	59%	58%
	Female	40%	38%	39%	40%	39%	40%	41%	42%
Percent by age group	0 to 19	14%	14%	13%	16%	14%	16%	16%	17%
	20 to 29	32%	29%	29%	27%	25%	23%	22%	19%
	30 to 39	24%	26%	27%	26%	28%	25%	25%	24%
	40 to 49	16%	16%	15%	14%	17%	19%	18%	19%
	50 to 59	11%	10%	11%	12%	11%	11%	12%	13%
	60 or more	4%	5%	5%	6%	6%	6%	7%	8%
	Cocaine	49%	52%	52%	46%	50%	49%	49%	46%
	Other psychostimulants (excl. cocaine)	62%	59%	58%	65%	62%	64%	65%	68%
Percent involving (categories not mutually exclusive)	Co-occurring non-stimulant poisonings	61%	62%	63%	60%	62%	61%	61%	65%
	Co-occurring opioid poisonings (incl. fentanyl or analogues)	40%	42%	42%	39%	42%	43%	43%	48%
	Co-occurring fentanyl or analogues poisonings	n/a	n/a	11%	14%	18%	24%	23%	28%
	Co-occurring non-stimulant/non-opioid poisonings	40%	38%	38%	35%	35%	33%	36%	36%
Median total length of stay (in days)		2	2	2	2	2	2	3	3

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
Totals used to calculate age-adjusted rates do not include cases with missing age.

The ICD-10-CA code for poisoning by fentanyl and derivatives was introduced April 2018. Fentanyl-specific data were not available prior to this date.
Data on opioid-related and stimulant-related poisonings hospitalizations are not mutually exclusive. A proportion of poisoning hospitalizations involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

Stimulant-related poisonings hospitalizations involving cocaine and other psychostimulants may exceed 100% as poisonings may co-occur.

Refer to the [Technical Notes](#) for more information.

Table 13c. Summary of intentional stimulant-related poisoning hospitalizations in Canada, 2016 to 2023 (January to June)

		2016	2017	2018	2019	2020	2021	2022	2023 (Jan to Jun)
Total	Number	649	699	696	703	729	698	657	329
	Crude rate	2.3	2.5	2.4	2.4	2.5	2.4	2.2	2.2
	Age adjusted rate	2.3	2.5	2.4	2.4	2.5	2.4	2.2	2.2
Percent by sex	Male	46%	49%	43%	44%	39%	39%	36%	37%
	Female	54%	51%	57%	56%	61%	61%	64%	63%
Percent by age group	0 to 19	25%	24%	27%	28%	30%	38%	37%	36%
	20 to 29	30%	29%	27%	27%	25%	25%	24%	27%
	30 to 39	21%	20%	21%	18%	21%	17%	19%	15%
	40 to 49	13%	15%	13%	12%	13%	11%	10%	12%
	50 to 59	9%	9%	7%	10%	7%	6%	7%	5%
	60 or more	2%	3%	5%	5%	4%	3%	3%	5%
	Cocaine	37%	40%	36%	33%	31%	29%	28%	26%
	Other psychostimulants (excl. cocaine)	69%	68%	70%	75%	75%	78%	80%	82%
Percent involving (categories not mutually exclusive)	Co-occurring non-stimulant poisonings	62%	65%	63%	61%	57%	51%	52%	57%
	Co-occurring opioid poisonings (incl. fentanyl or analogues)	35%	34%	32%	34%	30%	26%	26%	27%
	Co-occurring fentanyl or analogues poisonings	n/a	n/a	3%	5%	5%	8%	7%	8%
	Co-occurring non-stimulant/non-opioid poisonings	47%	48%	46%	41%	39%	35%	39%	41%
Not available (n/a)									

Note(s):

Data were not available at the time of this publication.

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Totals used to calculate age-adjusted rates do not include cases with missing age.
 The ICD-10-CA code for poisoning by fentanyl and derivatives was introduced April 2018. Fentanyl-specific data were not available prior to this date.
 Data on opioid-related and stimulant-related poisoning hospitalizations are not mutually exclusive. A proportion of poisoning hospitalizations involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.
 Refer to the [Technical Notes](#) for more information.

Table 14. Number of total stimulant-related poisoning hospitalizations by quarter and province or territory, January 2016 to June 2023

Province or territory	2016				2017				2018			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	123	107	146	104	123	152	141	167	113	146	142	147
AB	80	101	97	92	100	109	99	93	96	103	108	92
SK	29	34	28	29	21	29	28	30	23	37	33	33
MB	18	21	13	14	20	25	25	12	17	15	10	22
ON	138	206	202	176	189	245	247	235	251	302	299	255
QC	n/a											
NB	16	9	17	9	11	11	11	21	10	14	18	11
NS	Suppr.	Suppr.	11	11	10	8	10	9	Suppr.	9	Suppr.	8
PE	Suppr.	Suppr.	Suppr.	0	Suppr.	5	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NL	5	8	10	5	6	7	8	12	7	Suppr.	Suppr.	15
Territories	Suppr.	6	Suppr.	Suppr.	5	Suppr.						
Canada	409	492	524	440	485	586	574	579	517	626	610	583

Province or territory	2019				2020				2021			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	134	138	135	117	110	151	177	168	133	166	218	146
AB	72	81	86	82	78	131	110	106	87	115	96	104
SK	45	32	33	49	24	40	32	47	28	48	41	45
MB	13	14	11	20	6	17	20	13	11	18	12	13
ON	238	294	221	222	241	332	264	274	227	265	261	249
QC	n/a											
NB	16	12	16	12	14	14	19	12	15	15	14	16
NS	8	9	8	7	Suppr.	21	Suppr.	11	7	11	9	7
PE	0	Suppr.	Suppr.	Suppr.	0	0	Suppr.	Suppr.	0	Suppr.	0	Suppr.
NL	Suppr.	Suppr.	13	12	7	11	12	13	8	5	13	6
Territories	0	Suppr.	Suppr.	7	Suppr.	Suppr.	7	5	5	Suppr.	Suppr.	Suppr.
Canada	526	580	523	528	480	717	641	649	521	643	664	586

Province or territory	2022				2023	
	Q1	Q2	Q3	Q4	Q1	Q2
BC	155	163	187	168	150	155
AB	79	87	67	73	84	84
SK	28	26	24	38	40	35
MB	17	13	11	13	13	9
ON	217	203	235	189	228	254
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	12	16	17	16	13	6
NS	15	5	6	10	10	5
PE	Suppr.	Suppr.	0	Suppr.	Suppr.	0
NL	9	11	12	5	Suppr.	Suppr.
Territories	Suppr.	6	8	Suppr.	6	7
Canada	532	530	567	512	544	555

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Q1 = January to March; Q2 = April to June; Q3 = July to September; Q4 = October to December

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Totals for Canada by year may not match those from other tables as they do not include the suppressed numbers in provinces/territories, in order to comply with privacy and confidentiality rules.
 Refer to [Technical Notes](#) for more information.

Table 15. Number of accidental stimulant-related poisoning hospitalizations by quarter and province or territory, January 2016 to June 2023

Province or territory	2016				2017				2018			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	69	57	100	69	66	98	76	102	71	99	107	101
AB	39	54	51	47	54	58	62	54	53	61	54	55
SK	11	19	14	16	14	11	17	16	13	18	18	20
MB	10	10	8	5	8	16	16	6	7	10	Suppr.	Suppr.
ON	65	87	99	82	81	113	139	124	113	165	172	133
QC	n/a											
NB	5	Suppr.	0	0	Suppr.	Suppr.						
NS	Suppr.	0	Suppr.	5	Suppr.	Suppr.	Suppr.	Suppr.	0	0	5	0
PE	Suppr.	Suppr.	0	Suppr.	0	0	Suppr.	Suppr.	0	Suppr.	Suppr.	Suppr.
NL	0	Suppr.	Suppr.	0	Suppr.	Suppr.	0	Suppr.	0	Suppr.	Suppr.	Suppr.
Territories	Suppr.	Suppr.	0	0	Suppr.	Suppr.	0	Suppr.	0	Suppr.	Suppr.	Suppr.
Canada	199	227	272	224	223	296	310	302	257	358	351	309

Province or territory	2019				2020				2021			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
BC	85	69	78	56	58	100	121	96	85	106	149	85
AB	34	53	57	44	47	82	64	68	47	72	64	63
SK	16	17	10	20	8	18	17	29	12	20	24	22
MB	Suppr.	8	Suppr.	13	Suppr.	10	Suppr.	6	5	10	8	9
ON	123	174	122	131	136	203	163	168	125	154	153	144
QC	n/a											
NB	Suppr.	7	Suppr.	Suppr.	6	8	7	6	6	7	Suppr.	Suppr.
NS	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	5	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
PE	0	0	0	Suppr.	0	0	Suppr.	Suppr.	0	Suppr.	0	0
NL	0	Suppr.	Suppr.	0	Suppr.	Suppr.	Suppr.	0	Suppr.	Suppr.	Suppr.	Suppr.
Territories	0	Suppr.	0	Suppr.	Suppr.	Suppr.	Suppr.	0	Suppr.	Suppr.	Suppr.	Suppr.
Canada	258	328	267	264	255	426	372	373	280	369	398	323

Province or territory	2022				2023	
	Q1	Q2	Q3	Q4	Q1	Q1
BC	92	108	134	106	100	99
AB	52	52	33	32	54	47
SK	13	11	17	15	18	20
MB	11	5	Suppr.	Suppr.	Suppr.	Suppr.
ON	109	115	145	117	140	158
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	0	8	8	5	Suppr.	Suppr.
NS	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
PE	0	Suppr.	0	Suppr.	Suppr.	0
NL	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
Territories	0	Suppr.	6	Suppr.	Suppr.	Suppr.
Canada	277	299	343	275	312	324

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Q1 = January to March; Q2 = April to June; Q3 = July to September; Q4 = October to December

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.
 Totals for Canada by year may not match those from other tables, as they do not include the suppressed numbers in provinces/territories, in order to comply with privacy and confidentiality rules.
 Refer to [Technical Notes](#) for more information.

Table 16. Percentage of accidental stimulant-related poisoning hospitalizations by sex and province or territory, 2016 to 2023
(January to June)

Province or territory	2016		2017		2018		2019		2020		2021		2022		2023 (Jan to Jun)	
	Female	Male	Female	Male												
BC	29%	71%	25%	75%	28%	72%	31%	69%	24%	76%	31%	69%	28%	72%	27%	73%
AB	39%	61%	34%	66%	36%	64%	33%	67%	33%	67%	34%	66%	33%	67%	32%	68%
SK	35%	65%	43%	57%	38%	62%	52%	48%	39%	61%	44%	56%	41%	59%	50%	50%
MB	52%	48%	41%	59%	37%	63%	43%	57%	32%	68%	34%	66%	25%	75%	Suppr.	Suppr.
ON	32%	68%	30%	70%	31%	69%	32%	68%	29%	71%	29%	71%	28%	72%	30%	70%
QC	n/a	n/a														
NB	Suppr.	41%	59%	30%	70%	57%	43%									
NS	Suppr.	36%	64%	Suppr.	Suppr.	Suppr.	Suppr.									
PE	Suppr.	Suppr.														
NL	Suppr.	Suppr.														
Territories	Suppr.	Suppr.														
Canada	33%	67%	31%	69%	31%	69%	33%	67%	29%	71%	31%	69%	30%	70%	31%	69%

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

The data do not include cases with missing sex, or other sex.

Totals used to calculate Canada proportions do not include the suppressed numbers, in order to comply with privacy and confidentiality rules.

Refer to [Technical Notes](#) for more information.

Table 17. Percentage of accidental stimulant-related poisoning hospitalizations by age group and province or territory, 2016 to 2023 (January to June)

2016

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	11%	28%	23%	17%	15%	6%
AB	9%	40%	22%	15%	8%	7%
SK	Suppr.	27%	28%	20%	Suppr.	0%
MB	Suppr.	36%	15%	21%	15%	Suppr.
ON	5%	30%	31%	16%	13%	5%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	0%	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
NS	Suppr.	Suppr.	Suppr.	0%	Suppr.	0%
PE	0%	Suppr.	0%	Suppr.	0%	0%
NL	0%	0%	Suppr.	Suppr.	0%	Suppr.
Territories	0%	Suppr.	Suppr.	0%	0%	0%
Canada	8%	32%	26%	17%	12%	5%

2017

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	8%	29%	28%	19%	9%	6%
AB	11%	29%	29%	14%	10%	6%
SK	17%	36%	24%	Suppr.	12%	Suppr.
MB	11%	37%	22%	20%	Suppr.	Suppr.
ON	8%	26%	29%	18%	12%	6%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	Suppr.	Suppr.	0%	0%	0%
NS	Suppr.	Suppr.	Suppr.	Suppr.	0%	0%
PE	Suppr.	Suppr.	0%	0%	0%	0%
NL	0%	Suppr.	Suppr.	0%	0%	0%
Territories	0%	Suppr.	0%	Suppr.	0%	0%
Canada	9%	29%	29%	17%	11%	5%

2018

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	4%	30%	27%	17%	15%	7%
AB	7%	35%	26%	17%	8%	7%
SK	16%	33%	23%	16%	Suppr.	Suppr.
MB	Suppr.	26%	37%	19%	Suppr.	0%
ON	4%	28%	33%	16%	15%	4%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	0%	0%	Suppr.	Suppr.	0%	0%
NS	0%	Suppr.	Suppr.	Suppr.	0%	Suppr.
PE	Suppr.	0%	Suppr.	0%	0%	0%
NL	Suppr.	Suppr.	0%	0%	Suppr.	0%
Territories	0%	Suppr.	Suppr.	Suppr.	0%	0%
Canada	5%	30%	30%	17%	13%	5%

2019

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	11%	23%	31%	14%	13%	8%
AB	16%	24%	32%	14%	9%	5%
SK	17%	33%	25%	11%	Suppr.	Suppr.
MB	Suppr.	36%	39%	18%	0%	Suppr.
ON	6%	31%	28%	16%	14%	5%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	Suppr.	Suppr.	Suppr.	36%	0%
NS	Suppr.	Suppr.	50%	0%	0%	Suppr.
PE	Suppr.	0%	0%	0%	0%	Suppr.
NL	0%	Suppr.	Suppr.	0%	Suppr.	0%
Territories	Suppr.	0%	0%	0%	Suppr.	0%
Canada	10%	28%	30%	15%	12%	5%

2020

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	6%	21%	27%	21%	16%	10%
AB	8%	33%	32%	15%	9%	4%
SK	13%	29%	32%	19%	Suppr.	Suppr.
MB	Suppr.	28%	24%	32%	Suppr.	0%
ON	9%	24%	30%	18%	13%	7%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	19%	26%	22%	Suppr.	Suppr.
NS	0%	Suppr.	Suppr.	Suppr.	0%	Suppr.
PE	0%	Suppr.	Suppr.	0%	0%	0%
NL	Suppr.	0%	Suppr.	Suppr.	0%	0%
Territories	0%	0%	Suppr.	Suppr.	Suppr.	0%
Canada	8%	25%	30%	19%	12%	6%

2021

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	8%	16%	26%	21%	18%	11%
AB	11%	27%	27%	22%	7%	6%
SK	22%	31%	22%	15%	Suppr.	Suppr.
MB	Suppr.	34%	22%	Suppr.	16%	Suppr.
ON	5%	24%	29%	24%	11%	7%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	0%	Suppr.	Suppr.	40%	Suppr.	Suppr.
NS	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.	Suppr.
PE	Suppr.	0%	Suppr.	0%	0%	0%
NL	0%	Suppr.	Suppr.	Suppr.	Suppr.	0%
Territories	Suppr.	0%	Suppr.	0%	0%	0%
Canada	8%	23%	27%	22%	12%	7%

2022

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	5%	18%	25%	22%	16%	13%
AB	9%	25%	28%	21%	11%	5%
SK	16%	20%	30%	Suppr.	16%	Suppr.
MB	Suppr.	Suppr.	29%	21%	Suppr.	Suppr.
ON	7%	20%	30%	21%	15%	7%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	43%	Suppr.	Suppr.	24%	0%
NS	0%	Suppr.	Suppr.	Suppr.	0%	Suppr.
PE	Suppr.	0%	Suppr.	0%	0%	0%
NL	0%	Suppr.	Suppr.	Suppr.	0%	0%
Territories	0%	Suppr.	Suppr.	Suppr.	0%	Suppr.
Canada	7%	21%	28%	21%	15%	9%

2023 (Jan to Jun)

Province or territory	Age group					
	0 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 or more
BC	6%	14%	29%	22%	19%	11%
AB	11%	16%	31%	24%	14%	5%
SK	21%	32%	21%	18%	Suppr.	Suppr.
MB	Suppr.	Suppr.	Suppr.	0%	Suppr.	0%
ON	8%	15%	26%	23%	16%	12%
QC	n/a	n/a	n/a	n/a	n/a	n/a
NB	Suppr.	0%	Suppr.	Suppr.	Suppr.	Suppr.
NS	Suppr.	0%	Suppr.	0%	0%	Suppr.
PE	Suppr.	0%	0%	0%	0%	Suppr.
NL	Suppr.	0%	0%	Suppr.	Suppr.	Suppr.
Territories	0%	Suppr.	Suppr.	Suppr.	Suppr.	0%
Canada	9%	16%	28%	23%	16%	10%

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Not available (n/a) Data were not available at the time of this publication.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

The data do not include cases with missing age.

Totals used to calculate Canada proportions do not include the suppressed numbers, in order to comply with privacy and confidentiality rules.

Refer to [Technical Notes](#) for more information.

Table 18a. Number of total stimulant-related poisoning hospitalizations in Canada by sex and age group, 2016 to 2023 (January to June)

Age group by sex	2016		2017		2018		2019		2020		2021		2022		2023 (Jan to Jun)	
	Female	Male	Female	Male												
0 to 19	156	117	179	135	186	114	207	135	234	127	265	125	237	103	124	67
20 to 29	233	363	241	411	262	432	227	365	227	389	211	352	217	254	98	116
30 to 39	143	301	184	395	207	430	187	371	244	453	233	367	182	349	92	169
40 to 49	114	181	112	245	146	219	102	205	145	280	145	317	119	266	65	145
50 to 59	78	120	84	141	91	179	93	166	74	198	77	179	86	179	53	87
60 or more	33	45	38	68	43	64	56	64	44	98	39	109	48	107	29	64

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals for sex by age group may not match those from other tables as they do not include cases with missing sex, other sex or missing age.

Refer to [Technical Notes](#) for more information.

Table 18b. Number of intentional stimulant-related poisoning hospitalizations in Canada by age group and sex, 2016 to 2023 (January to June)

Age group by sex	2016		2017		2018		2019		2020		2021		2022		2023 (Jan to Jun)	
	Female	Male	Female	Male												
0 to 19	113	48	112	57	137	54	146	54	167	53	209	51	191	51	91	26
20 to 29	103	91	105	98	106	81	93	94	105	79	90	83	107	48	52	36
30 to 39	56	79	60	82	62	82	65	63	79	71	57	64	58	64	25	25
40 to 49	41	41	42	60	53	38	38	44	48	45	36	43	33	33	20	21
50 to 59	31	29	27	33	20	31	32	41	31	21	18	21	21	28	Suppr.	Suppr.
60 or more	8	8	10	12	16	16	21	12	13	15	11	12	12	10	7	9

Suppressed (Suppr.) Data may be suppressed in provinces or territories with low numbers of cases in order to comply with confidentiality rules.

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals for sex by age group may not match those from other tables as they do not include cases with missing sex, other sex or missing age.

Refer to [Technical Notes](#) for more information.

Table 18c. Number of accidental stimulant-related poisoning hospitalizations in Canada by age group and sex, 2016 to 2023 (January to June)

Age group by sex	2016		2017		2018		2019		2020		2021		2022		2023 (Jan to Jun)	
	Female	Male	Female	Male												
0 to 19	30	54	48	61	35	40	50	66	53	63	49	62	37	44	27	35
20 to 29	97	200	101	237	108	284	99	219	100	260	93	221	82	169	39	65
30 to 39	66	175	88	237	112	276	105	238	123	304	142	241	97	247	53	127
40 to 49	55	100	53	146	73	144	51	119	80	194	87	220	63	194	38	108
50 to 59	40	71	43	78	58	111	47	93	38	144	49	129	54	124	33	72
60 or more	22	30	21	42	22	42	26	38	27	69	24	84	29	81	19	48

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

Totals for sex by age group may not match those from other tables as they do not include cases with missing sex, other sex or missing age.

Refer to [Technical Notes](#) for more information.

Table 19. Percentage of accidental stimulant-related poisoning hospitalizations by substances involved, in Canada, 2016 to 2023
(January to June)

	2016	2017	2018	2019	2020	2021	2022	2023 (Jan to Jun)
Total number	932	1,145	1,298	1,147	1,449	1,402	1,216	657
Percent involving cocaine	55%	57%	59%	54%	58%	56%	59%	54%
Percent involving other psychostimulants (excl. cocaine)	59%	56%	53%	59%	57%	58%	58%	64%
Percent involving co-occurring non-stimulant poisonings	65%	63%	67%	61%	66%	67%	67%	70%
Percent involving co-occurring opioid poisonings (including fentanyl or fentanyl analogues)	45%	47%	49%	43%	49%	53%	54%	59%
Percent involving co-occurring fentanyl or fentanyl analogues poisoning	n/a	n/a	16%	20%	26%	32%	33%	38%
Percent involving co-occurring non-stimulant/non-opioid poisonings	39%	34%	36%	33%	34%	33%	36%	35%
Not available (n/a)	Data were not available at the time of this publication.							

Note(s):

Data on hospitalizations from Quebec are not included in the dataset used for analysis. As a result, Quebec is excluded.

The ICD-10-CA code for poisoning by fentanyl and derivatives was introduced April 2018. Fentanyl-specific data were not available prior to this date.

Data on opioid-related and stimulant-related poisoning hospitalizations are not mutually exclusive. A proportion of poisoning hospitalizations involving a stimulant also involved an opioid. Adding up those numbers would result in an overestimation of the burden of opioids and stimulants.

Stimulant-related poisoning hospitalizations involving cocaine and other psychostimulants may exceed 100% as poisonings may co-occur.

Refer to [Technical Notes](#) for more information.